Table 1: Characteristics of	people sample	e according to their n	mental health status (N=24,165)

Characteristics	Presence of any mental disorders	Absence of any mental
	% (95% CI)	disorder
	(n=2664)	% (95% CI) (n=21,501)
Age groups:		
15 - 24 years	29.2 (26.3, 32.1)	14.4 (13.7,15.2)
25 - 34 years	20.6 (17.3, 23.9)	15.5 (14.7,16.4)
35 - 44 years	16.8 (13.9, 19.7)	17.0 (16.1,18.0)
45 - 54 years	16.4 (13.5, 19.3)	18.7 (17.6,19.8)
55 - 64 years	11.6 (9.0, 34.0)	16.2 (15.4,17.0)
65 or more years	5.4 (42.4, 66.2)	18.2 (17.5,18.9)
Sex:		
Male	51.9 (47.6, 56.2)	49.0 (47.8,50.1)
Female	48.1 (43.8, 52.4)	51.0 (49.9,52.2)
Marital status:		
Married or Common law	40.8 (37.1, 44.5)	62.3 (61.1,63.4)
Widow or divorced or separated or single	59.1 (55.1, 63.1)	37.7 (36.6,38.9)
Education status:		
< Post-secondary Less than Post-secondary	47.0 (42.8, 51.2)	40.2 (39.1,41.4)
Post-secondary graduated	52.8 (48.4, 57.2)	59.8 (58.6,60.9)
Income status:		
No or less than \$20,000	8.6 (6.4, 10.8)	4.0 (3.6,4.4)
\$20,000 - \$39,999	15.4 (12.8, 18.0)	11.8 (11.1,12.4)
\$40,000 - \$59,999	18.1 (15.3, 20.9)	18.1 (17.2,19.0)
\$60,000 - \$79,999	17.3 (14.5, 20.1	17.6 (16.7,18.5)
\$80,000 or more	40.5 (36.9, 44.1)	48.6 (47.4,49.8)
Immigration:		
yes	14.1 (11.4, 16.8)	26.5 (15.3,27.6)
no	85.9 (81.9, 89.9)	73.5 (72.4,74.7)

Smoking status:			
Currently smoking	37.5 (34.0, 41.0)	19.1 (18.1,20.1)	
Former smoking	33.6 (29.8, 37.4)	39.3 (38.2,40.4)	
Never smoking	28.8 (25.432.2)	41.6 (40.5,42.8)	

CI: Confidence intervals corrected using the coefficient of variations provided by Statistics Canada

	With mental illness (n=2664)				Withou	t mental illness ((n=21,501)	
Characteristics	Currently smoking % (95% CI)	Former or quitted smoking % (95% CI)	Never smoking % (95% CI)	p- value	Currently smoking % (95% CI)	Former or quitted smoking % (95% CI)	Never smoking % (95% CI)	p- value
Age groups: 15 - 24 years	31.2 (26.4, 36.0)	21.1 (16.2, 26.0)	35.9 (30.1, 41.7)	<0.01	11.0 (9.6,12.6)	5.8 (5.1,6.5)	24.1 (22.7,25.6)	<0.01
25 - 34 years	23.0 (17.2, 28.8)	19.3 (13.5, 25.1)	18.9 (13.2, 24.6)		20.9 (18.6,23.3)	12.3 (11.2,13.5)	16.1 (14.7,17.7)	
35 - 44 years	16.7 (11.7, 21.7)	16.4 (11.4, 21.4)	17.1 (11.5, 22.7)		18.8 (16.4,21.4)	15.9 (14.5,17.5)	17.2 (15.8,18.7)	
45 - 54 years	16.1 (11.1, 21.1)	20.1 (14.1,26.1)	12.4 (7.5, 17.3) E		23.5 (20.7,26.5)	19.1 (17.5,20.8)	16.1 (14.5,18.0)	
55 - 64 years	10.5 (6.4, 14.6.) E	14.5 (9.5, 19.5) E	9.8 (5.3, 14.3) E		16.7 (14.9,18.6)	21.0 (19.6,22.5)	11.3 (10.3,12.4)	
65 or more years	2.4 (1.1, 3.7) E	8.4 (5.8, 11.0)	5.8 (3.5, 8.1) E		9.2 (8.1,10.3)	25.9 (24.6,27.2)	15.0 (14.0,16.1)	

Sex:								< 0.01
Male	54.2	56.8	43.3	< 0.01	57.6	54.3	39.9	
	(47.8, 60.6)	(50.1, 63.5)	(36.1, 50.5)		(54.7,60.5)	(52.5,56.1)	(38.1,41.8)	
Female	45.8	43.2	56.7		42.4	45.7	60.1	
	(39.6, 52.0)	(37.0, 49.4)	(49.7, 63.7)		(39.5,45.3)	(43.9,47.5)	(58.2,61.9)	
Marital status:								
Married or	33.7	50.7	38.6	< 0.01	57.9	71.6	55.5	< 0.01
Common law	(27.6, 39.8)	(44.4, 57.0)	(32.0, 45.2)		(55.0,60.7)	(70.0,73.1)	(53.7,57.4)	
Widow or	66.3	49.3	61.4		42.1	28.4	44.5	
divorced or separated or single	(58.5, 74.1)	(42.6, 56.0)	(53.8, 69.0)		(39.3,45.0)	(26.9,30.0)	(42.6,46.3)	
Education								
status:								
< Post-	59.6	37.7	42.0	< 0.01	45.8	35.9	41.8	< 0.01
secondary Less	(52.6, 66.6)	(31.2, 44.2)	(35.0, 49.0		(43.0,48.8)	(34.3,37.6)	(40.0,43.6)	
than Post- secondary								
Post-secondary	40.4	62.3	57.9		54.2	64.1	58.2	
graduated	(34.6, 46.2)	(54.9, 69.7)	(50.7, 65.1)		(51.2,57.0)	(62.4,65.7)	(56.4,60.0)	
Income status:								
no or < \$20,000	12.1	7.2	5.9	< 0.01	5.6	2.6	4.5	< 0.01
	(8.3, 15.9)	(4.0, 10.4) E	(2.7, 9.1) E		(4.8,6.5)	(2.2,3.2)	(3.8,5.3)	
\$20,000 -	19.3	11.8	14.4		13.7	11.3	11.3	
\$39,999	(13.9, 24.7)	(7.6, 16.0) E	(9.3, 19.5) E		(12.2,15.3)	(10.3,12.3)	(10.4,12.4)	
\$40,000 -	18.9	16.6	18.9		20.4	17.5	17.6	
\$59,999	(13.6, 24.2)	(11.8, 21.4)	(13.6, 24.2)		(18.2,22.8)	(16.3,18.7)	(16.2,19.0)	

\$60,000 -	13.8	22.2	16.1		17.2	17.4	17.9	
\$79,999	(9.5, 18.1)	(17.0, 27.4)	(11.2, 21.0)		(15.3,19.3)	(16.0,18.8)	(16.5,19.5)	
\$80,000 or more	35.9 (30.2, 41.6)	42.1 (35.6, 46.6)	44.7 (37.8, 51.6)		43.1 (40.1,46.2)	51.2 (49.5,53.0)	48.6 (46.8,50.5)	
Immigration:				< 0.01				
Yes	12.1 (7.8, 16.4) E	8.4 (4.9, 11.9) E	23.2 (23.2, 29.5)		19.0 (16.5,21.7)	20.4 (18.9,22.0)	35.7 (33.8,37.6)	< 0.01
No	87.9	91.6	76.9		81.0	79.6	64.3	
	(81.4, 94.4)	(87.8, 95.4)	(70.0, 83.8)		(78.3,83.5)	(78.0,81.1)	(62.4,66.2)	

Note: ^E: the coefficient of variation is in the marginal range, i.e. 16.6% - 33.3%. As suggested by Statistics Canada, the estimate should be interpreted with caution as there is high sampling variability associated with the estimate. CI: Confidence intervals corrected using the coefficient of variations provided by Statistics Canada p-value: chi2 p-value

Table 5: Association between characteristics of sample with their smoking status (n=2,664) among those with and without mental illness

Characteristics		h mental illness (former r restricted analysis)	Among people without mental illness (former vs. current smoker restricted analysis)		
	Unadjusted OR (95% CI)	Adjusted OR (95% CI)	Unadjusted OR (95% CI)	Adjusted OR (95% CI)	
Age groups:					
15 - 24 years	0.2 (0.1, 0.3)	0.2 (0.1, 0.4)	0.2 (0.1,0.2)	0.2 (0.2,0.3)	
25 - 34 years	0.2 (0.1, 0.4)	0.2 (0.1, 0.4)	0.2 (0.2,0.3)	0.2 (0.1,0.2)	
35 - 44 years	0.3 (0.2, 0.5)	0.2 (0.1, 0.4)	0.3 (0.2,0.4)	0.2 (0.2,0.3)	
45 - 54 years	0.4 (0.2, 0.7)	0.3 (0.1, 0.5)	0.3 (0.2,0.4)	0.2 (0.2,0.24)	
55 - 64 years	0.4(0.2, 0.7)	0.3 (0.2, 0.6)	0.4 (0.4,0.5)	0.3 (0.3,0.4)	
65 or more	Reference	Reference	Reference	Reference	
Sex:					
Female	0.9 (0.7, 1.2)	0.9 (0.6, 1.2)	1.1 (1.0,1.3)	1.2 (1.0,1.4)	
Male	Reference	Reference	Reference	1.3 Reference	
Marital status:					
Widow/divorced/separated or	0.5 (0.4, 0.7)	0.6 (0.4, 0.9)	0.5 (0.5,0.6)	0.7 (0.6,0.8)	
single	Reference	Reference	Reference	Reference	
Married/Common law					
Education status:					
<post-secondary< td=""><td>0.4 (0.3, 0.5)</td><td>0.4 (0.3, 0.6)</td><td>0.7 (0.6,0.8)</td><td>0.7 (0.6,0.8)</td></post-secondary<>	0.4 (0.3, 0.5)	0.4 (0.3, 0.6)	0.7 (0.6,0.8)	0.7 (0.6,0.8)	
Less than Post-secondary	Reference	Reference	Reference	Reference	
Post-secondary graduated					
Income status:					
no or less than \$20,000	0.5 (0.3, 0.8)	0.7 (0.4, 1.1)	0.4 (0.3,0.5)	0.4 (0.3,0.5)	
\$20,000 - \$39.999	0.5 (0.4, 0.8)	0.6 (0.4, 0.9)	0.7 (0.6,0.8)	0.5 (0.4,0.6)	
\$ 40.000 - \$ 59,999	0.7 (0.5, 1.1)	0.8 (0.5, 1.2)	0.7 (0.6,0.9)	0.6 (0.5,0.7)	
\$60,000 - \$ 79,999	1.4 (0.9, 2.1)	1.5 (1.0, 2.3)	0.8 (0.7,1.0)	0.8 (0.6,0.9)	

\$80,000 or more	Reference	Reference	Reference	Reference
Immigration:				
Yes	0.7 (0.4, 1.1)	0.6 (0.4, 0.9)	1.1 (0.9,1.3)	1.1 (0.9,1.3)
No	Reference	Reference	Reference	Reference

OR: odds ration; CI: confidence interval

Comparison of the odds ratios:

(With MI) Significant associations	(Without MI)
young and middle-aged	young and middle-aged
single or never married	single or never married
less educated	less educated (quitting more difficult among those with less edu, among MI)
immigrants	not significant
low income	low and middle income

The characteristics associated with smoking status (former vs. current smoking) were similar among those with and without mental disorders (refer to Table 5). Younger and middle age, single or never married status, less than post-secondary education, low income status and immigrant status were less likely to quit smoking, both among those with and without mental illness. However, compared to those without mental disorders, those with mental disorder were even more likely to continue smoking if they had less than post-secondary education. Immigrants were less likely to quit smoking if they had a mental illness, whereas there was no significant association between smoking and immigrant status among those without mental illness. *People without any mental illness belonging to middle income range were less likely to quit smoking whereas this was not the case among those with mental illness.*

This finding reinforces that people with less than post-secondary education and immigrant status are at high risk of smoking cessation problems if they are suffering from any kind of mental illness and interventions have to be directed towards them while implementing smoking cessation programs.

Table 6: Physical and mental health characteristics of people across smoking status among those with mental illness (N=24,165)

	With			
Characteristics	Currently smoking % (95% CI)	Former or quitted smoking % (95% CI)	Never smoking % (95% CI)	p-value
Physical activity Yes No	71.2 (67.2,74.9) 28.8 (25.1,32.8)	74.3 (69.8,78.4) 25.7 (21.6,30.2)	74.6 (69.0,79.4) 25.4 (20.6,31.0)	0.4
Chronic condition At least one None	58.5 (53.7,63.1) 41.5 (36.9,46.3)	59.3 (54.2,64.2) 40.7 (35.8,45.8)	56.9 (50.8,62.7) 43.1 (37.3,49.2)	0.8
Professional mental health consultation Yes No	43.3 (38.7,48.1) 56.7 (51.9,61.3)	45.4 (40.5,50.4) 54.6 (49.6,59.5)	47.4 (41.2,53.7) 52.6 (46.3,58.8)	0.5

© 2020 Dahar R. et al.