

Supplementary Table 1. Demographic characteristics of study participants and available quit smoking supports in their clinic (n=40)

| Variable | n | % |
|---|-------------|----------|
| Sex | | |
| Male | 9 | 22.5 |
| Female | 31 | 77.5 |
| Age | | |
| <29 years | 4 | 10.0 |
| 30–39 years | 12 | 30.0 |
| 40–49 years | 9 | 22.5 |
| 50–59 years | 12 | 30.0 |
| 60–69 years | 3 | 7.5 |
| Country | | |
| Armenia | 8 | 20.0 |
| Georgia | 9 | 22.5 |
| Greece | 2 | 5.0 |
| Romania | 8 | 20.0 |
| Russia | 6 | 15.0 |
| Ukraine | 7 | 17.5 |
| Primary practice setting | | |
| Solo | 2 | 5.0 |
| Group | 3 | 7.5 |
| Hospital | 16 | 40.0 |
| Primary Care | 4 | 10.0 |
| Community | 7 | 17.5 |
| Other (NGO) | 8 | 20.0 |
| Area of practice | | |
| Urban | 39 | 97.5 |
| Rural | 1 | 2.5 |
| Average patients per week, mean (SD) | 29.7 (30.1) | |
| Years-of-practice, mean (SD) | 21.4 (24.2) | |
| Employment status | | |
| Part-time | 8 | 20.0 |
| Full-time | 32 | 80.0 |
| Previous participation in a smoking cessation program | | |
| Yes | 15 | 37.5 |
| No | 25 | 62.5 |
| Personal experience with the tobacco use | | |
| Former smoker | 8 | 20.0 |
| Non-smoker | 32 | 80.0 |
| Which of the following supports are available in your clinic? | | |
| Process to screen and document smoking status of patients | 22 | 55.0 |
| Self-help materials of smokers | 20 | 50.0 |
| Consult forms to guide you through quit smoking counselling interventions | 12 | 30.0 |
| None of the above | 8 | 20.0 |
| Other supports | 8 | 20.0 |